



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

From (d) and (e),  $\Psi\delta\psi + \Phi\delta\phi + \Theta\delta\theta = -\delta V \dots (f)$ .

$$\therefore \Psi = -\frac{dV}{d\psi}, \quad \Phi = -\frac{dV}{d\phi}, \quad \text{and} \quad \Theta = -\frac{dV}{d\theta}.$$

Therefore the Lagrangian equations of motion may be written:

$$\frac{d}{dt}\left(dT \Big/ \frac{d\psi}{dt}\right) = \frac{dT}{d\psi} - \frac{dV}{d\psi}, \quad \frac{d}{dt}\left(dT \Big/ \frac{d\phi}{dt}\right) = \frac{dT}{d\phi} - \frac{dV}{d\phi},$$

$$\text{and} \quad \frac{d}{dt}\left(dT \Big/ \frac{d\theta}{dt}\right) = \frac{dT}{d\theta} - \frac{dV}{d\theta}.$$

## NON-EUCLIDEAN GEOMETRY: HISTORICAL AND EXPOSITORY.

BY GEORGE BRUCE HALSTED, A. M., (Princeton) Ph. D., (Johns Hopkins); Member of the London Mathematical Society; and Professor of Mathematics in the University of Texas, Austin, Texas.

[Continued from the February Number].

**PROPOSITION XIV.** *The hypothesis of obtuse angle is inconsistent with Euclid's assumption: Two straight lines cannot enclose a space.*


**Proof.** From the hypothesis of obtuse angle, assumed as true, [and the first 28 propositions of Euclid], we have now deduced the truth of Euclid's Postulatum; that two straights will meet each other in some point toward those parts, toward which a certain straight, cutting them, makes two internal angles, of whatever kind, less than two right angles.

But this Postulatum holding good, on which Euclid supports himself after the twenty-eighth proposition of his first book, it is manifest to all Geometers, that the hypothesis of right angle alone is true, nor any place left for the hypothesis of obtuse angle. Therefore the hypothesis of obtuse angle is inconsistent with Euclid's assumption.

Quod erat demonstrandum.

Otherwise, and more immediately.

Since from the hypothesis of obtuse angle we have demonstrated (P. IX.) that two (fig. 11.) acute angles of the triangle  $APX$ , right-angled at  $P$ , are greater than one right angle; it follows that an acute angle  $PAD$  may be assumed such, that together with the aforesaid two acute angles it makes up two right angles. But then the straight  $AD$  must (by the preceding proposition,



joined to the hypothesis of obtuse angle) at length meet with this  $PL$ , or  $XL$ , regard being had to the secant, or incident  $AP$ ; which is manifestly absurd against Eu. I. 17, if we regard the secant or incident  $AX$ .

**PROPOSITION XV.** *By any triangle  $ABC$ , of which the three angles (fig. 13.) are together equal to, or greater, or less than two right angles, is established respectively the hypothesis of right angle, or obtuse angle, or acute angle.*

**Proof.** For anyhow two angles of this triangle, as suppose  $A$ , and  $C$ , will be acute, because of Eu. I. 17. Wherefore the perpendicular, let fall from the apex of the remaining angle  $B$  upon this  $AC$ , will cut this  $AC$  (Eu. I. 17.) in some intermediate point  $D$ .

If therefore three angles of this triangle  $ABC$  are supposed equal to two right angles, it follows that all the angles of the triangles  $ADB$ ,  $CDB$  will be together equal to four right angles, because of the two additional right angles at the point  $D$ . This holding good, now of neither of the said triangles, as suppose  $ADB$ , will the three angles together be less, or greater than two right angles; for thus viceversa the three angles together of the other triangle would be greater, or less than two right angles. Wherefore (P. IX.) from one triangle indeed would be established the hypothesis of acute angle, and from the other the hypothesis of obtuse angle; which is contrary to P. VI. and P. VII.

Therefore the three angles together of either of the aforesaid triangles will be equal to two right angles; and therefore (P. IX.) is established the hypothesis of right angle.

Quod erat primo loco demonstrandum.

But if however the three angles of the proposed triangle  $ABC$  are taken greater than two right angles; now of the two triangles  $ADB$ ,  $CDB$  all the angles together will be greater than four right angles, because of the two additional right angles at the point  $D$ .


This holding good; now of neither of the said triangles will the three angles together be precisely equal to, or less than two right angles; for thus viceversa the three angles of the other triangle would be together greater than two right angles. Wherefore (P. IX.) from one triangle indeed would be established the hypothesis either of right angle or of acute angle, and from the other the hypothesis of obtuse angle, which is contrary to P. V, P. VI, and P. VII.

Therefore the three angles together of either of the aforesaid triangles will be greater than two right angles; and therefore (P. IX.) is established the hypothesis of obtuse angle.

Quod erat secundo loco demonstrandum.

But finally. If the three angles of the proposed triangle  $ABC$  are taken less than two right angles, now of the two triangles  $ADB$ ,  $CDB$ , all the angles together will be less than four right angles, because of the two additional right angles at the point  $D$ .

This holding good; now of neither of the said triangles will the three angles together be equal to, or greater than two right angles; for thus viceversa of the other triangle the three angles together would be less than two right angles. Wherefore (P. IX.) from one triangle indeed would be establish-



ed the hypothesis either of right angle or obtuse angle, and from the other the hypothesis of acute angle; which is contrary to P.V, P.VI, and P.VII.

Therefore the three angles together of either of the aforesaid triangles will be less than two right angles; and therefore (P.IX.) is established the hypothesis of acute angle. *Quod erat tertio loco demonstrandum.*

Accordingly by any triangle  $ABC$ , of which the three angles are equal to, or greater, or less than two right angles, is established respectively the hypothesis of right angle, or obtuse angle, or acute angle. *Quod erat propositum.*

**COROLLARY.** Hence, any one side of any proposed triangle being produced, as suppose  $AB$  to  $H$ , the external angle  $HBC$  will be (Eu.I.13.) either equal to, or less, or greater than the remaining internal and opposite angles together at the points  $A$  and  $C$ , according as is true the hypothesis of right angle, or obtuse angle, or acute angle. And inversely.

[To be continued.]

---

## THE "IRREDUCIBLE CASE."

---

By J. K. ELLWOOD, A. M., Principal Colfax School, Pittsburg, Pa.

---

**PROBLEM.**—To extract the cube root of  $a \pm \sqrt{-b}$ .

Put  $\sqrt[3]{a + \sqrt{-b}} = m + n$ , and  $\sqrt[3]{a - \sqrt{-b}} = m - n$ .

Then  $a + \sqrt{-b} = m^3 + 3m^2n + 3mn^2 + n^3$ , and  $a - \sqrt{-b} = m^3 - 3m^2n + 3mn^2 - n^3$ . Hence  $a = m^3 + 3mn^2$ , and  $\sqrt{-b} = 3m^2n + n^3$ .

*Example 1.* Find the cube root of  $9 + 25\sqrt{-2}$ .

Here  $a = m^3 + 3mn^2 = 9 = 3^3 - 18$ . Hence  $3mn^2 = -18$ , and  $n = \sqrt{-2}$ .

To verify these values of  $m$  and  $n$  substitute them in  $\sqrt{-b} = 3m^2n + n^3$   $= 25\sqrt{-2}$ . Doing this we have  $27\sqrt{-2} + (-2\sqrt{-2}) = 25\sqrt{-2}$ .

$\therefore 3 + \sqrt{-2}$  is the required root. When the substituted values of  $m$  and  $n$  do not give the second term they are not correct, and other values must be found by trial.

*Example 2.* Find the cube root of  $2\sqrt{11} + 30\sqrt{-3}$ . Here  $a = m^3 + 3mn^2 = 2\sqrt{11} = (\sqrt{11})^3 - 9\sqrt{11}$ .

Hence  $3mn^2 = -9\sqrt{11}$ , and  $n = \sqrt{-3}$ . Since these values of  $m$  and  $n$  substituted in  $3m^2n + n^3$  give  $\sqrt{-b} = 30\sqrt{-3}$ , the root is  $\sqrt{11} + \sqrt{-3}$ .

This method frequently enables us to simplify Cardan's formula for cubics in what is called the "irreducible case", said formula being

$$x = \sqrt[3]{q + \sqrt{(q^2 - p^3)}} + \sqrt[3]{q - \sqrt{(q^2 - p^3)}}.$$